

## ग्रामोभ्युदयादेव देशोभ्युदयः


Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



### Gramin Krishi Mausam Sewa Bulletin, District - Udham Singh Nagar

Year: 28 Issue No.: 1 Bulletin Period: 02– 06 January, 2019 Day: Tuesday Dated: 01-01-2019

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Udham Singh Nagar and plain area of Nainital districts are as follows:

Parameter	Udham Singh Nagar				
	02/01/2019	03/01/2019	04/01/2019	05/01/2019	06/01/2019
Rainfall (mm)	0	0	0	0	2
Max Temp ( <sup>o</sup> C)	22	23	22	21	22
Min Temp ( <sup>o</sup> C)	4	3	3	2	4
Cloud Cover	P-Cloudy	P-Cloudy	Clear	Cloudy	Cloudy
Max RH I (%)	85	90	85	90	85
Min RH II (%)	40	45	40	45	40
Wind speed (km/h)	004	004	004	006	004
Wind direction	WNW	ESE	NW	WNW	ESE

According to Meteorological observations recorded at Agrometeorological Observatory, AMFU-Pantnagar, G B Pant University of Agriculture & Technology (Altitude: 243.8 m) during last seven days (25-31, December 2018, 2018) are as follows:

Weather Parameter	Udham Singh Nagar						
	25/12/2018	26/12/2018	27/12/2018	28/12/2018	29/12/2018	30/12/2018	31/12/2018
Rainfall (mm)	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Max Temp ( <sup>o</sup> C)	22.0	20.5	19.0	19.7	18.5	19.0	21.0
Min Temp ( <sup>o</sup> C)	4.4	2.4	2.9	1.8	0.9	2.8	0.6
CC I	4	4	4	4	4	4	4
CC II	0	4	4	0	0	4	4
Max RH I (%)	97	100	100	100	100	90	96
Min RH II (%)	59	43	62	42	41	62	53
Wind speed (km/h)	NR	NR	NR	NR	NR	NR	NR
Wind direction I	N	Clear	Clear	N	Calm	Calm	N
Wind Direction II	WSW	WSW	WSW	WNW	ESE	WNW	WSW

## Weather Based Agro-Advisories

### Crop Management:

- If there is highly occurrence of broad leaf weed in the wheat crop, then solution of 500g AI of 2,4D or 4g of metsulfuron methyl or 20g of carfentrazone in 500-600 liter of water should be sprayed @ per hectare with the help of flat fan nozzle after 34-40 days in November sown wheat crop and after 40-45 days in December sown wheat crop.
- On the highly occurrence of narrow leaf weed in the wheat crop, Pinoxaden @ 40-45g/Ha or Sulfosulfuron @ 25g/ha or Clodinafop 60g/Ha or Finoxaprop Ethyl @ 100-120g/Ha should be sprayed after 30-35 days of sowing of wheat crop.
- If there is a mixed occurrence of broad and narrow leaf weed, then Sulfosulfuron + Metsulfuron (30+2) @ 32g/500-600 liter of water should be sprayed within 25-30days of sowing.
- It is advised that wheat crop should not be sown in the month of January because wheat sowing during this period causes the yield reduction of 60-65Kg/Ha/day and hot wind during ripening stage of crop causes thinning of wheat grain.
- First irrigation should be given at an interval of 20-25 days of sowing in the standing wheat crop at root initiation stange and second irrigation should be given at 40-45 days at the sprouting of wheat.
- To control the leaf spot disease in mustard, solution of mancozeb @ 2.5 g/liter of water should be sprayed.


### Horticultural Management:

- Top dressing of Urea should be done in Cabbage and Cauliflower crop after weeding and hoeing and moisture should be maintained in the field.
- Potato leaves should be cut near 10<sup>th</sup> January.
- To control the late blight disease in potato and tomato, it is advised that solution of Mancozeb @ 2.5g/liter or copper oxichloride @ 3g/lit of water should be sprayed.
- On the withering of brinjal, solution of carbendazim @1 g/ lit of water should be sprayed. If whole plant is drying then root should be irrigated with the same solution.
- When lower leaves of pea are turning yellow and plants are drying, then roots of pea crop should be irrigated with the solution of carbendazim @ 1 g/ lit of water.
- To control the leaf spot disease in cole crops, solution of mancozeb @ 2.5 g/liter of water should be sprayed.
- Irrigation should be given in the new orchards to save them from frost.
- To control the Mango mealy bug in mango, polythene strips should be used so that plants can be protected from these insects. For this, stem of plants should be covered by 25-30 cm broad polythene and above 30-40 cm from the ground. After covering the stem, grease and unusable oil should be applied in the upper and lower part of polythene and both the end of polythene should be tied with the help of chord/string.

### Animal Husbandry:

- To save the animals from cold, proper arrangement of animal shed should be done.
- Sitting place of animals should be leveled so that their productive capacity is not affected and special care should be given to the new born animals.
- To save the animal from cold, dry grasses, paddy residue (Puwal) etc which are not used as fodder for animals should be used as a bedding material for animals in sheds. Door and window should be covered properly so that cold wind could not be entered in the animal shed.
- To save the animal from cold, increase the amount of oil and Jaggery (Gur) in the food of animals. Caraway (Ajwain) and gur also be given to the animals.

- In hilly areas, heater may be used to protect the animals from cold. Rice husk should be given to the animals as a food which is the source of energy.
- Maintain the temperature of dwelling place of poultry birds.
- With the changing weather, there is a high chance of Niomonina in new born animals. So, it is advisable that animal shed should be protected from cold and hot food should be given to the animals.
- Keeping in mind the probable high rate of Cattle's delivery, animal shed should be well neat, clean, dry and ventilated. For this, dry Lime should be sprayed in the nearby places.
- Toxocara vitulorum (patera) are found in the food tube of buffalo calf in the age of 1-4month. More than 50% buffalo calf should be died because of delay in treatment. Piperazine should be given to the infected calf on identification of Patera disease.
- To save the buffalo calf from patera disease, 10-15Cc Neem oil should be given to the calf immediately after 10 days of his birth. Again after 10 days, 10-15 CC Neem oil should be given. Chenopodium album oil can also be given as a fodder to animals to save the calf/animals from patera disease.
- It is advised that green fodder can be given to the animals by mixing it in dry fodder otherwise animal may be infected by Timpeti disease which causes death of animals.
- There is a chance of Aflatoxicosis disease in Poultry due to occurrence of fungus in the food which causes to death of poultry birds. So, medicine should be administered to the poultry birds only on the advice of veterinarian.



**Dr. R K Singh**  
**Professor & Principal Nodal Officer-GKMS**  
**AMFU- Pantnagar**