

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Nainital

Year: 28 Issues No.: 18 Bulletin Periods: 2 - 6 March, 2019 Day: Friday Dated: 01-03-2019

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Nainital district are as follows:


Parameter	Nainital				
	02/03/2019	03/03/2019	04/03/2019	05/03/2019	06/03/2019
Rainfall (mm)	0	5	3	3	3
Max Temp (⁰ C)	12	14	11	12	12
Min Temp (⁰ C)	2	3	3	2	2
Cloud Cover	P-Cloudy	Dense Cloudy	Cloudy	Cloudy	Cloudy
Max RH I (%)	80	85	85	85	85
Min RH II (%)	40	45	45	45	45
Wind speed (km/h)	006	004	006	004	004
Wind direction	NW	NW	ESE	NNW	NNW

Light rain may likely to occur from 3rd to 6th of March with cloudy to dense cloudy sky.

According to the observations recorded at India Meteorological Department Observatory located at Nainital (Altitude: 2084 m) during last seven days (22-28 Feb, 2019), it is observed that state of sky were cloudy to dense cloudy. Maximum and minimum temperature varied between 6.7⁰C to 16.6⁰C and 1.9⁰C to 5.3⁰C respectively.

Weather Based Agro-Advisories

Crop	Stage	Agromet Advisory
Wheat, Barley	Booting	Keeping in view the rainfall condition in coming days, farmers are advised to stop irrigation for next few days in all the crops.
Sugarcane	Early-Harvesting Late-Sowing	Harvesting of early sown sugarcane crop should be done. Sowing of spring season sugarcane crop should be completed up to 15 th March. Upper 2/3 part of sugarcane should be used for sugarcane seed. 40-50 thousand pieces of three eyed sugarcane should be used. Line should be prepared at a distance of 75cm in East-West direction for sowing sugarcane crop.
Onion	Vegetative Growth	Weeding and hoeing should be done.
Tomato	Early-Plucking Late-Weeding and hoeing	Plucking of tomato fruit should be done in early sown variety of tomato. Weeding and hoeing should be done in late sown tomato crop.
Brinjal	-	Weeding and hoeing should be done.
Apple, pear, plum, wall nut etc	Sowing	Sowing of fruit crops like-apple, pear, plum, wall nut etc should be done.
Animal Husbandry	-	Keeping in mind the probable high rate of Cattle's delivery, animal shed should be well neat, clean, dry and ventilated. For this, dry Lime should be sprayed in the nearby places.
Poultry	-	Maintain the temperature of dwelling place of poultry birds. To save the poultry bird from cold, thickness of bedding material should be increases so that they can get sufficient heat to maintain their temperature.



Dr. R K Singh
Professor & Principal Nodal Officer-GKMS
AMFU- Pantnagar