

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Udham Singh Nagar

Year: 27 Issue No.: 35 Bulletin Period: 2 – 6 May, 2018 Day: Tuesday Dated: 01-05-2018

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Udham Singh Nagar and plain area of Nainital districts are as follows:

Parameter	Udham Singh Nagar				
	02/05/2018	03/05/2018	04/05/2018	05/05/2018	06/05/2018
Rainfall (mm)	0	15	5	0	0
Max Temp (⁰ C)	35	33	31	33	34
Min Temp (⁰ C)	19	19	18	18	19
Cloud Cover	P-Cloudy	Dense Cloudy	Dense Cloudy	Cloudy	P-Cloudy
Max RH I (%)	80	85	90	85	80
Min RH II (%)	40	45	50	45	45
Wind speed (km/h)	006	008	006	004	004
Wind direction	SE	E	SE	SE	NE

Moderate rain on 3rd May & light rain on 4th May with dense cloudy sky may likely to occur.

According to Meteorological observations recorded at Agrometeorological Observatory, AMFU-Pantnagar, G B Pant University of Agriculture & Technology (Altitude: 243.8 m) during last seven days (24 – 30 April, 2018) are as follows:

Weather Parameter	Udham Singh Nagar						
	24/04/2018	25/04/2018	26/04/2018	27/04/2018	28/04/2018	29/04/2018	30/04/2018
Rainfall (mm)	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Max Temp (⁰ C)	36.0	37.4	38.5	36.5	34.5	35.0	35.5
Min Temp (⁰ C)	16.3	17.1	29.4	25.9	21.4	21.5	21.0
CC I	Clear	Clear	Cloudy	P-Cloudy	Cloudy	P-Cloudy	Clear
CC II	Clear	Clear	Cloudy	P-Cloudy	Clear	P-Cloudy	Clear
Max RH I (%)	80	58	60	61	60	62	65
Min RH II (%)	18	31	40	48	40	52	31
Wind speed (km/h)	4.0	3.1	4.8	13.9	9.7	7.2	9.8
Wind direction I	Calm	Calm	WSW	ESE	ESE	ESE	SSE
Wind Direction II	WNW	WNW	ESE	ENE	SSE	ESE	ESE

Weather Based Agro-Advisories

Crop Management:


- Bunding and summer ploughing of the field should be done in this month.
- If irrigation facility is available then sowing of dhaincha and sanai should be done after harvesting of wheat and other rabi crops for green manure.
- Progressive variety of sanai like Narendra sanai-1 and dhaincha variety like pant dhaincha-1, hisar dhaincha-1 etc should be selected. Seed rate for sanai should be 80-90 kg/ha and for dhaincha 60-80 kg/ha.
- Soil sample should be taken from different places in this month from fallow field after harvesting of rabi crops. Soil sample should be taken up to the depth of 20 cm.
- In paddy-wheat crop rotation, after harvesting of wheat or any other rabi crops, sowing of green manure like sesbania (dhaincha) and sannhemp (sanai) should be done up to 15th May.
- To control the root and stem rot diseases in french bean and lobia, carbendazim @ 1g/lit of water should be sprayed.

Horticultural Management:

- Transplanting of capsicum and brinjal should be done in the polyhouse in mid and high hills and required irrigation should be applied.
- On the occurrence of angular spot in the leaves of cucurbitaceous (pumpkin class vegetables) crop, solution of mancozeb @ 2.5 kg/lit of water should be sprayed.
- To control the leaf blight disease in onion, tebuconazole or difinoconazole or propiconazole @ 500ml/ha with any srvangi insecticide and sticker should be sprayed.
- On the occurrence of contracted piebald leaves in chilly and tomato, infected plants should be removed and destroyed. To control the juice sucking insect in these crops, sarvangi insecticide should be sprayed. To save the crop from late blight disease, it is advised that solution of Mancozeb @ 2.5g/liter or copper oxichloride @ 3g/lit of water should be sprayed.
- To control fruit fly in mango orchard, pheromone trap made up of wood should be should be placed on tree (10 trap/ha) and pheromone trap should be changed in 2 months.
- To control the black tip and internal necrosis, 1.0 % borax (10g/liter) should be sprayed.
- Necessary irrigation should be applied to maintain the proper moisture condition in the orchard.
- To control the powdery mildew in mango, solution of 0.1 % carbendazim should be sprayed.

Animal Husbandry:

- In the summer season, time to time spraying of melathiyani or other insecticide should be done in the drains near the animal shed.
- Temperature of animal shed should be maintained using cooling devices such as fan, cooler or latest cooling device to maintain the productivity of foreign cows and save them from diseases.
- When animal is affected by sun stroke then contact the veterinarian immediately for their remedy.
- Neat, fresh and cold water should be given to the animals three times in a day. Availability of sufficient moisture in the body of animals maintains the balance between the temperature of their skin and environment which save the animals from the effect of loo (summer wind).
- To save the animals from high temperature, balanced diet should be given to them. Green fodder mixed with dry fodder and grain must be given to the animals.
- Buffalo should be kept in the animal shed from 9 AM to 6 PM, to save them from sun.

A handwritten signature in black ink, appearing to read 'R K Singh', is enclosed in a light purple rectangular box.

Dr. R K Singh
Professor & Principal Nodal Officer-GKMS
AMFU- Pantnagar