

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Udham Singh Nagar

Year: 26 Issue No.: 35 Bulletin Period: 3-7 May, 2017 Day: Tuesday Dated: 02-05-2017

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Udham Singh Nagar and plain area of Nainital districts are as follows:

Parameter	Udham Singh Nagar				
	03/05/2017	04/05/2017	05/05/2017	06/05/2017	07/05/2017
Rainfall (mm)	0	0	0	0	0
Max Temp (⁰ C)	35	35	34	35	35
Min Temp (⁰ C)	17	19	20	19	19
Cloud Cover	Cloudy	Cloudy	Cloudy	Cloudy	P-Cloudy
Max RH I (%)	80	80	85	85	85
Min RH II (%)	40	40	50	50	50
Wind speed (km/h)	08	08	12	08	08
Wind direction	ESE	SE	NNW	ESE	SE

According to Meteorological observations recorded at Agrometeorological Observatory, AMFU- Pantnagar, G B Pant University of Agriculture & Technology (Altitude: 243.8 m) during last seven days (25 April to 01 May, 2017) are as follows:

Weather Parameter	Udham Singh Nagar						
	25/04/2017	26/04/2017	27/04/2017	28/04/2017	29/04/2017	30/04/2017	01/05/2017
Rainfall (mm)	0.0	0.0	0.0	0.0	0.0	0.0	Trace
Max Temp (⁰ C)	38.6	38.5	37.6	37.0	38.5	39.6	37.0
Min Temp (⁰ C)	18.9	22.4	19.4	18.4	20.1	19.9	20.4
CC I	Clear	Cloudy	Clear	Clear	Clear	Dense Cloud	Dense Cloud
CC II	Clear	Cloudy	Cloudy	Cloudy	P-Cloudy	Cloudy	Clear
Max RH I (%)	70	55	63	46	52	69	54
Min RH II (%)	21	22	18	17	19	22	31
Wind speed (km/h)	7.3	13.4	8.7	7.7	8.3	7.6	8.0
Wind direction I	WNW	WNW	Clear	NNE	WNW	S	ESE
Wind Direction II	WNW	WSW	WNW	WNW	SSW	ENE	ENE

Weather Based Agro-Advisories

Crop Management:

- If irrigation facility is available then sowing of dhaincha and sanai should be done after harvesting of wheat and other rabi crops for green manure.
- Progressive variety of sanai like Narendra sanai-1 and dhaincha variety like pant dhaincha-1, hisar dhaincha-1 etc should be selected. Seed rate for sanai should be 80-90 kg/ha and for dhaincha 60-80 kg/ha.
- Soil sample should be taken from different places in this month from fallow field after harvesting of rabi crops. Soil sample should be taken up to the depth of 20 cm.
- Harvesting and threshing of wheat and pulse crops should be completed.

Horticultural Management:

- To control dansi makkhi in mango orchard, pheromone trap made up of wood should be placed on tree (10 trap/ha) and pheromone trap should be changed in 2 months.
- To control the internal necrosis or koyaliya, 0.8-1.0 % (10g/liter) borax should be sprayed.
- Necessary irrigation should be applied to maintain the proper moisture condition in the orchard.
- On the occurrence of yellow spot of irregular shape on cucurbitaceous crops, leaves should be examined by turning and if there is a growth of light gray fungus in the lower portion of leaves then it should be controlled by spraying the solution of Mancozeb @ 2.5 Kg/ Liter.
- If the veins of leaves of okra crop are turning yellow, then destroy the plants and Sarvangi insecticide should be sprayed to control the juice sucking insects.
- To control the powdery mildew in mango, solution of 0.1 % carbendazim should be sprayed.
- On the occurrence of fine mottled spot in upper leaves of chilly and tomato, Sarvangi insecticide at 10-15 days interval should be sprayed.
- On the occurrence of fruit borer in the Tomato, fruit should be used only after 3 days and 5 days of spraying of Chlorantraniliprole 18.5 SC @ 150ml/Ha and Indoxacarb 14.5 SC @ 500ml/Ha respectively.
- To control the stem and fruit borer in brinjal, spraying of Emamectin benzoate 5SG @ 200g/Ha, Cypermethrin 25EC @200ml/Ha, Lambda cyhalothrin 5CS @ 300ml/Ha should be done and fruit should be used after 5 days of last spraying of chemicals.

Animal Husbandry:

- In the summer season, time to time spraying of melathiyani or other insecticide should be done in the drains near the animal shed.
- Temperature of animal shed should be maintained using cooling devices such as fan, cooler or latest cooling device to maintain the productivity of foreign cows and save them from diseases.
- When animal is affected by sun stroke then contact the veterinarian immediately for their remedy.
- Neat, fresh and cold water should be given to the animals three times in a day. Availability of sufficient moisture in the body of animals maintains the balance between the temperature of their skin and environment which save the animals from the effect of loo (summer wind).
- To save the animals from high temperature, balanced diet should be given to them. Green fodder mixed with dry fodder and grain must be given to the animals.
- Buffalo should be kept in the animal shed from 9 AM to 6 PM, to save them from sun.

Dr. R K Singh
Professor & Nodal Officer-GKMS
AMFU- Pantnagar