

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Udham Singh Nagar

Year: 26 Issue No.: 78 Bulletin Period: 7 – 11 Oct, 2017 Day: Friday Dated: 06-10-2017

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Udham Singh Nagar and plain area of Nainital districts are as follows:

Parameter	Udham Singh Nagar				
	07/10/2017	08/10/2017	09/10/2017	10/10/2017	11/10/2017
Rainfall (mm)	0	0	0	0	0
Max Temp (⁰ C)	34	33	33	32	32
Min Temp (⁰ C)	23	23	22	22	23
Cloud Cover	P-Cloudy	Clear	P-Cloudy	P-Cloudy	Cloudy
Max RH I (%)	85	85	85	90	90
Min RH II (%)	45	45	45	50	50
Wind speed (km/h)	004	006	006	006	008
Wind direction	WNW	NW	SE	ESE	ESE

According to Meteorological observations recorded at Agrometeorological Observatory, AMFU-Pantnagar, G B Pant University of Agriculture & Technology (Altitude: 243.8 m) during last seven days (29 Sept - 05 Oct, 2017) are as follows:

Weather Parameter	Udham Singh Nagar						
	29/09/2017	30/09/2017	01/10/2017	02/10/2017	03/10/2017	04/10/2017	05/10/2017
Rainfall (mm)	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Max Temp (⁰ C)	33.5	34.0	32.8	33.0	32.5	32.5	33.6
Min Temp (⁰ C)	23.5	24.0	24.4	24.5	24.5	24.0	23.9
CC I CC II	Dense Cloud	Dense Cloud	Overcast	Clear	Dense Cloud	Dense Cloud	P-Cloudy
	Clear	Overcast	Dense Cloud	Dense Cloud	Dense Cloud	Dense Cloud	Cloudy
Max RH I (%)	88	81	88	85	90	87	88
Min RH II (%)	57	73	62	64	65	65	66
Wind speed (km/h)	1.4	1.3	3.2	3.1	3.0	2.5	0.9
Wind direction I	ENE	ENE	ENE	ENE	ENE	N	Calm
Wind Direction II	WNW	ESE	WSW	WSW	WSW	WSW	ESE

Weather Based Agro-Advisories

Crop Management:

- If field will remain empty after first week of October, then Rai and Mustard (toria) should be sown in first fortnight of October.
- For sowing of Rai and Mustard crop, Varieties of Rai like- Varuna, Rohini, Krishna, Kanti, Vardan, Vaibhav, Basanti, Narendra, Early rai-4, Urvashi and varieties of Yellow Mustard like- Ragini, Binoy (B-9), Pant peeli Sarso-1 should be selected.
- If the cover of corn turn yellow in the Maize then picking of corn should be done. After picking of corn, green plants of Maize should be used as a fodder for animals.
- In winter season variety of Sugarcane crop, sowing of should be completed up to 15 October.
- Harvesting of early and mid season variety of Paddy should be done on the appearance of golden color of panicle in Paddy.
- Irrigation should be stopped before 10-15 days of harvesting of paddy. For this, moisture level in the grains reaches up to 20% and 10-15cm irrigation should be saved.
- To control the bacterial leaf blight disease in paddy crop, streptomycin 15 g + copper oxichloride 500 g / hectare should be sprayed.
- If growth of Sugarcane is good then tie the plants of three basins of the two rows by using Kaichi method.
- On the occurrence of yellow mosaic disease in the Pulse (Dalhani) crops, leaves turn yellow. This disease is spread by the White Fly. To control this, 2-3 spraying of 30 EC Dimethoate or 20 EC Methyl-O-dimeton @ 1liter/Ha in 500-600 liter of water should be done at an interval of 10-12 days.
- On the attack of stem borer, Chlorantraniliprole 20 SC @ 150ml/ha or Kartop 50SP @ 1kg/Ha or Flubendiamide 480 SC @75ml/Ha or Monocrotophos 36 SL @ 1400 ml/Ha should be sprayed.

Horticultural Management:

- On the occurrence of termite, Imidacloprid @ 1ml/liter or Chlorpyrifos @ 4ml/liter should be use in the basin of the stem of fruit trees.
- Clearing and ploughing of orchard should be done.
- Paclobutrazol should be used for proper fruiting of trees.
- In case of web on the trees, cleaning of web may be done by using web machine and Quinalphos @ 2ml/liter should be sprayed.
- Transplanting of late variety of Cauliflower can also be done up to the second and third week of this month. 150 Kg N, 80 Kg P, 60 Kg K doses of fertilizers are required. From these doses, half dose of Nitrogen and full dose of Phosphorous should be applied at the time of final ploughing and the remaining dose of Nitrogen should be given at 30-35 days and 60-65 days after transplanting of plants.
- If the fruits of early variety of Cauliflower were ready in the fields then send them tom the market and top dressing of Urea should be given in the mid variety of Cauliflower sown in the last month. Arrangements of weeding, hoeing and irrigation should be done.
- If there is an occurrence of leaf sucking insects (Mahu & White Fly), solution of 0.1% concentration of Metasystox or Imidachorpid should be sprayed and Sandovit or powder of soap should be mixed in the solution.
- On the occurrence of leaf spot disease in brinjal crop, solution of mancozeb @ 2g/lit of water should be sprayed.

- To control the sucking insect in the Colocasia, 0.2 % solution of Imidachlorpid + Dithene 45 should be sprayed on 10 days interval.
- To save the Cauliflower from attack of insect, 0.1 % Metasystox or Imidachlorpid should be sprayed.

Animal Husbandry:

- In green fodder, leguminous fodder is best diet for animals and to increase production. So it is advised that cattleman should grow leguminous fodder (berseem) to keep their cattle healthy and to obtain maximum production from them.
- In this month, rate of animal's delivery especially in buffalos increases. The pregnant animals should be separated from other animals if possible and complete diet in little quantity at many times in a day should be given to them otherwise they can be affected by afra (fatty liver).
- Just after delivery, tie the navel of new born baby of cattle by cutting it with sharp and clean knife, then apply Gentian violet paint or tincture iodine.
- Within 2 hours of animal's delivery, little quantity of colostrums should be given to the new born baby of cattle after their proper cleaning.
- Green fodder should be given in less quantity to the animals. It is advised that green fodder can be given to the animals by mixing it in dry fodder.
- Drinking water should be clean because there is a possibility of occurrence of viruses due to parasite and fungus.
- Animal shed should be dried, for this; lime should be sprayed time to time in their places.

Dr. R K Singh
Professor & Principal Nodal Officer-GKMS
AMFU- Pantnagar