

# ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



## Gramin Krishi Mausam Sewa Bulletin, District - Nainital

Year: 27 Issue No.: 12 Bulletin Period: 10 – 14 Feb 2018 Day: Friday Dated: 09.02.2018

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Nainital district are as follows:

Parameter	Nainital				
	10/02/2018	11/02/2018	12/02/2018	13/02/2018	14/02/2018
Rainfall (mm)	0	0	5	15	2
Max Temp ( $^{\circ}$ C)	13	14	13	10	10
Min Temp ( $^{\circ}$ C)	3	4	5	3	2
Cloud Cover	P-Cloudy	Cloudy	Cloudy	Dense Cloudy	Cloudy
Max RH I (%)	80	80	85	85	85
Min RH II (%)	45	40	45	45	45
Wind speed (km/h)	008	008	006	006	006
Wind direction	NW	NW	SW	SE	ESE

**Due to western disturbances, light to moderate rain with high wind may likely to occur from 12<sup>th</sup> to 14<sup>th</sup> February 2018 with cloudy to dense cloudy sky.**

According to the observations recorded at India Meteorological Department Observatory located at Nainital (Altitude: 2084 m) during last seven days (2–8 February 2018), it is observed that state of sky were clear to cloudy and rainfall of 0.0 mm was received. Maximum and minimum temperature varied between 12.4 $^{\circ}$ C to 17.4 $^{\circ}$ C and 3.3 $^{\circ}$ C to 6.5 $^{\circ}$ C respectively.

### Weather Based Agro-Advisories

#### Crop Management:

- Weather will remain dry. On the occurrence of yellow blisters in the lower leaves or brown spot in the leaves or withering of leaves by turning yellow from the tip in the wheat crop, Propiconazole 25EC @ 1lit/ha should be sprayed.
- Weeding, hoeing, irrigation and nutrients should be applied in the wheat crop.
- Preparation for sowing spring season sugarcane crop should be done.
- Second irrigation should be given to the wheat crop after 40-45 days of sowing. Spraying of remaining 1/3<sup>rd</sup> dose of urea should be done.
- Advance agricultural equipments should be used to control the weed in the wheat crop.

- Irrigation should be applied in the pulse crops before pod formation and weed control should be done.
- Harvesting of Naulakh sugarcane crop should be done in mid February for good ratoon of sugarcane crop.
- On the occurrence of Mahu insect in the wheat crop, Thiamethoxam 25WSG @50g/Ha or Quinolphos 25 EC @1liter/Ha should be sprayed.
- Harvesting of matured crop of Mustard (Toria) should be done.

### **Horticultural Management:**


- On the occurrence of leaf minor insect in pea crop, 0.08 % solution of Imidachloprid should be sprayed.
- In valley, field should be prepared for sowing of potato in the last week of this month. 4-5 quintal dung manure with 2.5 kg DAP and 2 kg Potash fertilizer should be used.
- Light irrigation should be given after harvesting of the vegetable crops, sown in the valleys.
- When lower leaves of pea are turning yellow and plants are drying, then roots of pea crop should be irrigated with the solution of carbendazim @ 1 g/ lit of water.
- If the upper part of leaves of onion and garlic are turning yellow, solution of propiconazole or tebuconazole @ 1 ml/lit of water should be sprayed.
- To control the late blight disease in potato and tomato, it is advised that solution of Mancozeb @ 2.5g/liter or copper oxichloride @ 3g/lit of water should be sprayed.
- On the occurrence of yellowish brown spot in tomato, mancozeb 2.5-3.0 g/lit of water should be sprayed.
- To control the leaf spot disease in cole crops, solution of mancozeb @ 2.5 g/liter of water should be sprayed.
- Ensure the nutrient supply in the temperate fruit trees in mid and high hills.
- To control the leaf curl disease in the seed bearing fruit like Peach, plum etc recommended insecticide should be used.
- Tongue grafting should be started in the nursery for high quality temperate fruit plant.
- Sowing of fruit crops like-apple, pear, plum, wall nut etc should be started.
- Basin should be prepared in the temperate fruit plants and right quantity of dung manure, Nitrogen and Phosphorous should be applied in the field.
- To control the Canker disease in the apple, 0.3% Copper-oxichloride should be used after cutting and pruning. Unwanted and affected branches of apple should be removed through pruning and insecticide and fungicide should be sprayed.
- To control the fruit Rot disease in apple and seed bearing fruits, soil should be removed from the surrounded area of stem in the affected fruit tree so that sun rays may enter directly in the stem of affected tree. Affected bark should be removed and chaubatia paste should be applied and covered with soil. Drenching of 0.3 % Copper-oxichloride/plant should also be done.
- In high hills, pruning of temperate fruit trees like apple, pear, peach, plum etc should be done.

### **Animal Husbandry:**

- Maintain the temperature of dwelling place of poultry birds. To save the poultry bird from cold, thickness of bedding material should be increases so that they can get sufficient heat to maintain their temperature.
- To save the animals from cold, proper arrangement of animal shed should be done. To save the animal from cold, dry grasses, paddy residue (Puwal) etc which are not used as fodder for animals

should be used as a bedding material for animals in sheds. Door and window should be covered properly so that cold wind could not be entered in the animal shed.

- Sitting place of animals should be leveled so that their productive capacity is not affected and special care should be given to the new born animals.
- To save the animal from cold, increase the amount of oil and Jaggery (Gur) in the food of animals. Caraway (Ajwain) and gur also be given to the animals.
- Keeping in mind the probable high rate of Cattle's delivery, animal shed should be well neat, clean, dry and ventilated. For this, dry Lime should be sprayed in the nearby places.
- Toxocara vitulorum (patera) are found in the food tube of buffalo calf in the age of 1-4month. More than 50% buffalo calf should be died because of delay in treatment. Piperazine should be given to the infected calf on identification of Patera disease.
- To save the buffalo calf from patera disease, 10-15Cc Neem oil should be given to the calf immediately after 10 days of his birth. Again after 10 days, 10-15 CC Neem oil should be given. Chenopodium album oil can also be given as a fodder to animals to save the calf/animals from patera disease.



**Dr. R K Singh**

**Professor & Principal Nodal Officer-GKMS  
AMFU- Pantnagar**