

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Udham Singh Nagar

Year: 27 Issue No.: 20 Bulletin Period: 10 – 14 March, 2018 Day: Friday Dated: 09-03-2018

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Udham Singh Nagar and plain area of Nainital districts are as follows:

Parameter	Udham Singh Nagar				
	10/03/2018	11/03/2018	12/03/2018	13/03/2018	14/03/2018
Rainfall (mm)	0	0	0	0	0
Max Temp (⁰ C)	30	30	30	31	31
Min Temp (⁰ C)	10	10	11	10	10
Cloud Cover	P-Cloudy	P-Cloudy	P-Cloudy	P-Cloudy	P-Cloudy
Max RH I (%)	85	85	80	80	80
Min RH II (%)	40	40	40	40	40
Wind speed (km/h)	004	008	008	006	004
Wind direction	NW	NW	WNW	WNW	WNW

According to Meteorological observations recorded at Agrometeorological Observatory, AMFU-Pantnagar, G B Pant University of Agriculture & Technology (Altitude: 243.8 m) during last seven days (2 March – 8 March, 2018) are as follows:

Weather Parameter	Udham Singh Nagar						
	02/03/2018	03/03/2018	04/03/2018	05/03/2018	06/03/2018	07/03/2018	08/03/2018
Rainfall (mm)	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Max Temp (⁰ C)	28.4	30.0	30.5	31.8	28.2	28.0	28.5
Min Temp (⁰ C)	10.5	12.0	12.4	12.0	10.9	9.9	11.0
CC I CC II	Cloudy	Cloudy	Cloudy	P-Cloudy	Clear	P-Cloudy	P-Cloudy
	Clear	Dense Cloudy	Dense Cloudy	P-Cloudy	Clear	Clear	Clear
Max RH I (%)	95	91	81	90	90	95	90
Min RH II (%)	40	29	36	42	42	53	39
Wind speed (km/h)	5.0	6.4	6.2	5.4	5.4	7.4	2.8
Wind direction I	Calm	WNW	WNW	Calm	Calm	Calm	Calm
Wind Direction II	WSW	WSW	WSW	WNW	WSW	WNW	WSW

Weather Based Agro-Advisories

Crop Management:


- Immediate supply of late sown variety of sugarcane to the mills should be done after harvesting. Urea should be applied in the field and cover the field with 10cm thick layer of dry leaves before taking ratoon crop. If more than 45cm space is found in the lines then fill them with the pieces of sugarcane. More than 20% Nitrogen should be used in ratoon crop as compare to the Naulakh sugarcane crop. Half dose of nitrogen should be given at the time of first irrigation and remaining half dose at the second and third irrigation.
- Sowing of spring season sugarcane crop should be completed till 15th March. Seeds should be taken from the registered seeds nursery. Seed should be treated in the solution of Carbendazim for 10 minutes before sowing.
- Sowing of Urd should be till 15th March completed.
- Sowing of sunflower and groundnut variety should be completed upto 15th March.
- Modern hybrid variety of sunflower like BSH-1, Arun, Paras, Divyamukhi and jwalamukhi should be selected for sowing. 10-12 Kg/Ha seeds should be sown in lines at a distance of 45 cm and plant to plant distance would be 20cm. 80 Kg Nitrogen, 60Kg Phosphorous and 40Kg potash per Hectare should be applied.
- Ground nut variety like Punjab Moongfali-1, Type-64, Kaushal, Prakash, Amber etc should be sown in lines at a distance of 30cm with plant to plant distance of 10cm. Seed rate should be 80-100Kg/Ha. 20 Kg Nitrogen, 40Kg Phosphorous and 45 Kg potash /Ha should be used.
- Fodder crops like chari, Maize, bajara, makchari, lobia and jawar etc can also be sown in this month.
- On the occurrence of brown color, circular ring shape spot in the mustard class crop, mancozeb @ 1.5-2.0 kg/lit should be sprayed.
- On the occurrence of yellow blisters in the lower leaves or brown spot in the leaves or withering of leaves by turning yellow from the tip in the wheat crop, Propiconazole 25EC @ 1lit/ha should be sprayed.
- On the occurrence of yellow gerui in wheat crop, Propiconazole 25 EC (or tilt) @ 500ml/Ha in 500 liter of water should be sprayed.

Horticultural Management:

- Field preparation should be done for sowing bottle gourd (lauki), bitter gourd (Karela), ridged gourd (Torai) and pumpkin (Kashiphal) crop.
- When lower leaves of pea are turning yellow and plants are drying, then roots of pea crop should be irrigated with the solution of carbendazim @ 1 g/ lit of water.
- If the upper part of leaves of onion and garlic are turning yellow, solution of propiconazole or tebuconazole @ 1 ml/lit of water should be sprayed.
- To control the late blight disease in potato and tomato, it is advised that solution of Mancozeb @ 2.5g/liter or copper oxichloride @ 3g/lit of water should be sprayed.
- On the occurrence of yellowish brown spot in tomato, mancozeb 2.5-3.0 g/lit of water should be sprayed.
- To control the leaf spot disease in cole crops, solution of mancozeb @ 2.5 g/liter of water should be sprayed.
- To control the mango hopper (Bhunga) insect in fruit trees, first spray should be given at panicle initiation with Imidacloprid 17.8 SL @ 0.03 ml/lit. Second spray at pea stage with thiamethoxam @ 0.32 g/lit and third spray need based at 21 days after second spray with NSKE 5% @ 5ml/lit.

Animal Husbandry:

- To prevent Puerperal Fever in Pregnant animals, 50-60 grams of mineral mixture should be fed to them every day to boost their immunity.
- In this month care needs to be taken to prevent disease caused by hot water.
- If mosquitoes, flies, ticks etc. are increasing due care needs to be taken to prevent spread of disease caused by them.
- Animals are prone to sterility and Johne's disease (Paratuberculosis) during this time. They should provided immediate treatment for these.
- To save the animals from cold, proper arrangement of animal shed should be done. To save the animal from cold, dry grasses, paddy residue (Puwal) etc which are not used as fodder for animals should be used as a bedding material for animals in sheds.
- Sitting place of animals should be leveled so that their productive capacity is not affected and special care should be given to the new born animals.
- To save the animal from cold, increase the amount of oil and Jaggery (Gur) in the food of animals. Caraway (Ajwain) and gur also be given to the animals.
- Keeping in mind the probable high rate of Cattle's delivery, animal shed should be well neat, clean, dry and ventilated. For this, dry Lime should be sprayed in the nearby places.
- Toxocara vitulorum (patera) are found in the food tube of buffalo calf in the age of 1-4month. More than 50% buffalo calf should be died because of delay in treatment. Piperazine should be given to the infected calf on identification of Patera disease.
- To save the buffalo calf from patera disease, 10-15Cc Neem oil should be given to the calf immediately after 10 days of his birth. Again after 10 days, 10-15 CC Neem oil should be given. Chenopodium album oil can also be given as a fodder to animals to save the calf/animals from patera disease.



Dr. R K Singh
Professor & Principal Nodal Officer-GKMS
AMFU- Pantnagar