

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Udham Singh Nagar

Year: 26 Issue No.: 44 Bulletin Period: 10 - 14 June, 2017 Day: Friday Dated: 09-06-2017

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Udham Singh Nagar and plain area of Nainital districts are as follows:

Parameter	Udham Singh Nagar				
	10/06/2017	11/06/2017	12/06/2017	13/06/2017	14/06/2017
Rainfall (mm)	0	0	0	0	0
Max Temp (⁰ C)	35	37	38	38	39
Min Temp (⁰ C)	23	24	24	25	26
Cloud Cover	Cloudy	P-Cloudy	P-Cloudy	P-Cloudy	P-Cloudy
Max RH I (%)	85	85	80	80	80
Min RH II (%)	45	45	40	40	40
Wind speed (km/h)	06	08	08	10	10
Wind direction	NW	ESE	NE	NW	ESE

According to Meteorological observations recorded at Agrometeorological Observatory, AMFU- Pantnagar, G B Pant University of Agriculture & Technology (Altitude: 243.8 m) during last seven days (2 - 8 June, 2017) are as follows:

Weather Parameter	Udham Singh Nagar						
	02/06/2017	03/06/2017	04/06/2017	05/06/2017	06/06/2017	07/06/2017	08/06/2017
Rainfall (mm)	0.0	0.0	0.0	0.0	0.0	7.0	0.0
Max Temp (⁰ C)	38.5	40.5	42.0	42.0	42.5	34.0	32.0
Min Temp (⁰ C)	22.0	22.4	27.0	29.6	28.0	24.5	22.5
CC I	Clear	Clear	Cloudy	Cloudy	P-Cloudy	Dense Cloud	Clear
CC II	Clear	Cloudy	Cloudy	Cloudy	P-Cloudy	P-Cloudy	Clear
Max RH I (%)	62	59	54	62	58	78	70
Min RH II (%)	21	23	33	32	55	80	53
Wind speed (km/h)	5.8	7.8	9.1	8.5	11.2	13.9	5.2
Wind direction I	ENE	ENE	ESE	ENE	ENE	NNE	ENE
Wind Direction II	WSW	WNW	ENE	WNW	ESE	ESE	WSW

Weather Based Agro-Advisories

Crop Management:

- Harvest the urd and moong crops immediately after maturing. Don't delay in the harvesting otherwise grains will fall after cracking of pods. Plucking of pods should be done after maturing of 80% pods of urd and moong and plough the green plants in the field as green manure.
- Nursery of paddy should be prepared.
- Direct sowing of paddy should be done in the first fortnight of June and for this mid season variety should be selected. Sowing should be done in lines at a distance of 18-22cm with the seed rate of 60-70Kg/Ha. 1Kg Active Ingredient of pendimethalin should be sprayed within 2 days of sowing. It saves 20% water and the expenditure of transplanting and crops are ready before 7-10 days.
- Sowing of maize should be done in the first fortnight of June. Seed rate should be 18-20 kg/ha for sowing suitable varieties like Naveen, Tarun, Kiran, Navjot, Pratap Makka-1, Pusa sankul 3 and 4. Sowing should be done at a distance of 60-75cm. After germination plant to plant distance should be kept 20-25 cm through thinning. After sowing 40-50 kg nitrogen, 60 kg P₂O₅ and 40 kg/ha potash should be used.
- Sowing of early maturing variety of pigeon pea (arhar) should be done in the first fortnight of June so that late sowing of wheat can be done after that.
- Variety of cowpea like pant arhar-3, Pant arhar-291, UPS-120, Pusa-992 etc should be sown at a distance of 50-60cm in lines and seed rate should be 15-20 kg/ha. Seed should be treated with rhizobium and 15 Kg nitrogen and 40-50 Kg Phosphorous should be used at the time of sowing.
- On the occurrence of black bug in sugarcane crop, Chlorpyrifos 20 EC @ 2liter/Ha or phenthoate 50EC @ 1 liter/ha or 2 liter of quinalphos 25 EC in 500 liter of water should be sprayed.

Horticultural Management:

- To control the dryness of the liana in cucurbitaceous crops, destroy the dry liana after cutting them and roots of crops should be irrigated with the solution of 0.1% Carbendazim (1g/liter).
- If the upper stem of chilly crop is drying after turning black, then infected branches should be removed by plucking to save the crop and solution of 0.1% Carbendazim should be sprayed.
- On the occurrence of yellow spot of irregular shape on cucurbitaceous crops, leaves should be examined by turning and if there is a growth of light gray fungus in the lower portion of leaves then it should be controlled by spraying the solution of Mancozeb @ 2.5 Kg/ Liter.
- If the veins of leaves of okra crop are turning yellow, then destroy the plants and Sarvangi insecticide should be sprayed to control the juice sucking insects.
- On the occurrence of fine mottled spot in upper leaves of chilly and tomato, Sarvangi insecticide at 10-15 days interval should be sprayed.
- On the attack of White fly in tomato, Cyantraniliprole 10.26 OD @ 900ml/Ha or Thiamethoxam 25 WSG @ 200g/Ha should be applied and fruit should be used after 5 days of spraying.
- To save the chilly from Thrips, Lambda cyhalothrin 5EC @300ml/Ha or Fipronil 5SC @ 1Liter/Ha should be sprayed and chilly is used after 7 days of spraying of chemicals.
- To control the Mite in chilly, diafenthiuron 50WP @ 600g/Ha or Lambda cyhalothrin 5EC @ 300ml/Ha should be sprayed and in this case fruit may be used after 5 days of spraying.
- On the occurrence of fruit borer in the Tomato, fruit should be used only after 3 days and 5 days of spraying of Chlorantraniliprole 18.5 SC @ 150ml/Ha and Indoxacarb 14.5 SC @ 500ml/Ha respectively.
- To control the stem and fruit borer in brinjal, spraying of Emamectin benzoate 5SG @ 200g/Ha, Cypermethrin 25EC @200ml/Ha, Lambda cyhalothrin 5CS @ 300ml/Ha should be done and fruit should be used after 5 days of last spraying of chemicals.

Animal Husbandry:

- To save the animals from infected disease, vaccination should be done on the recommendation of veterinarian.
- On the occurrence of nitrate toxicity in the food of animals, respiration and pulse rate of animals increases and they feel weakness and stiffness in their muscles. Give the 50-100 ml dose of 1% Methylene blue direct in the veins of animals if their food is affected by nitrogen toxicity.
- If animal has consumed the cyanide affected fodder then water should not be given to them. Animals should be kept away from grazing the less grown jawar, bajara and fodder in pasture.
- Small shrivel/fade yellow stiff plants should not be used as fodder for animals to save them from cyanide.
- In the summer season, time to time spraying of melathiyam or other insecticide should be done in the drains near the animal shed.
- Temperature of animal shed should be maintained using cooling devices such as fan, cooler or latest cooling device to maintain the productivity of foreign cows and save them from diseases.
- When animal is affected by sun stroke then contact the veterinarian immediately for their remedy.
- Neat, fresh and cold water should be given to the animals three times in a day. Availability of sufficient moisture in the body of animals maintains the balance between the temperature of their skin and environment which save the animals from the effect of loo (summer wind).
- To save the animals from high temperature, balanced diet should be given to them. Green fodder mixed with dry fodder and grain must be given to the animals.

Dr. R K Singh
Professor & Nodal Officer-GKMS
AMFU- Pantnagar