

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Nainital

Year: 27 Issue No.: 46 Bulletin Period: 13-17 June 2018 Day: Tuesday Dated: 12.06.2018

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Nainital district are as follows:

Parameter	Nainital				
	13/06/2018	14/06/2018	15/06/2018	16/06/2018	17/06/2018
Rainfall (mm)	1	0	1	3	5
Max Temp ($^{\circ}$ C)	26	27	27	26	25
Min Temp ($^{\circ}$ C)	14	14	13	14	14
Cloud Cover	Partly cloudy	Clear	Partly Cloudy	cloudy	cloudy
Max RH I (%)	80	80	80	90	90
Min RH II (%)	40	40	40	45	50
Wind speed (km/h)	006	008	004	010	012
Wind direction	ESE	SE	WNW	WNW	WNW

According to the observations recorded at India Meteorological Department Observatory located at Nainital (Altitude: 2084 m) during last seven days (05-11 June, 2018), it is observed that state of sky were p-cloudy, cloudy to dense cloudy. Maximum and minimum temperature varied between 22.0° C to 24.0° C and 12.0° C to 13° C respectively.

Weather Based Agro-Advisories

Crop Management:


- Bunding of the field should be done for nursery preparation of paddy.
- Preparation of nursery of paddy should be completed up to first week of June.
- Pure seeds of paddy should be used for nursery i.e. weeds should not be mixed in the seeds.
- Arrangements for safety of crops should be done to save them from insect-pest attack.
- In spring season sugarcane, if broad leaves weed are there then 2, 4 D-Sodium salt @ 0.75-1.0 Kg AI/ha dissolve in 750 liter of water should be sprayed in 1 ha area. If agriculture labour are available then hoeing should be done and leave them for 1 week after that irrigation should be done and apply Nitrogen in the form of Urea in the presence of sufficient moisture.
- If the leaves of french bean and lobia are drying, carbendazim @ 1 g/lit of water should be used.

Horticultural Management:

- Continue the removal of shoots emerges from root stock of the grafting plants in the nursery.
- To control the gummosis disease in fruit bearing fruit trees, Streptocycline 0.01% or copper oxichloride 0.025% should be sprayed at 15 days interval.
- In mid hilly regions, to save the apple from rotting, plenofix @ 10ppm should be sprayed.
- Picking of early variety of peach, plum and khumani should be done and send them to the market.
- In high hilly regions, to control the scab disease on the falling of pankhuri of flower in apple crop, Carbendazim 0.05 % should be used.
- In valley, irrigation should be stop at the maturity stage of crops like rai, radish, turnip, cauliflower and harvesting should be done at the maturity of crops.
- On the occurrence of spot on the leaves of tomato and chilly, Mancozeb @ 2.5g/liter of water should be sprayed.
- Transplanting of capsicum and brinjal should be done in the polyhouse in mid and high hills and required irrigation should be applied.
- To control the leaf blight disease in onion, tebuconazole or difinoconazole or propiconazole @ 500ml/ha with any sarvangi insecticide and sticker should be sprayed.
- To control To prevent leaf blot and fruit rot disease in tomatoes, make a molecule 2.5 grams per liter of water and make a spraying solution
- On the occurrence of contracted piebald leaves in chilly and tomato, infected plants should be removed and destroyed. To control the juice sucking insect in these crops, sarvangi insecticide should be sprayed. To save the crop from late blight disease, it is advised that solution of Mancozeb @ 2.5g/liter or copper oxichloride @ 3g/lit of water should be sprayed.

Animal Husbandry:

- On the occurrence of nitrate toxicity in the food of animals, respiration and pulse rate of animals increases and they feel weakness and stiffness in their muscles. Give the 50-100 ml dose of 1% Methylene blue direct in the veins of animals if their food is affected by nitrogen toxicity.
- If animal has consumed the cyanide affected fodder then water should not be given to them. Animals should be kept away from grazing of less grown jawar, bajara, fodder in pasture.
- Small shrivel/fade yellow stiff plants should not be used as fodder for animals to save them from cyanide.
- In the summer season, time to time spraying of melathiyan or other insecticide should be done in the drains near the animal shed.
- Temperature of animal shed should be maintained using cooling devices such as fan, cooler or latest cooling device to maintain the productivity of foreign cows and save them from diseases.
- When animal is affected by sun stroke then contact the veterinarian immediately for their remedy.
- Neat, fresh and cold water should be given to the animals three times in a day. Availability of sufficient moisture in the body of animals maintains the balance between the temperature of their skin and environment which save the animals from the effect of loo (summer wind).
- To save the animals from high temperature, balanced diet should be given to the animals. Green fodder mixed with dry fodder and grain must be given to the animals.



Dr. R K Singh

**Professor & Principal Nodal Officer-GKMS
AMFU- Pantnagar**