

# ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



## Gramin Krishi Mausam Sewa Bulletin, District - Udham Singh Nagar

Year: 26 Issue No.: 66 Bulletin Period: 26 - 30 August, 2017 Day: Friday Dated: 25-08-2017

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Udham Singh Nagar and plain area of Nainital districts are as follows:

Parameter	Udham Singh Nagar				
	26/08/2017	27/08/2017	28/08/2017	29/08/2017	30/08/2017
Rainfall (mm)	15	20	20	05	10
Max Temp ( <sup>0</sup> C)	32	31	31	32	33
Min Temp ( <sup>0</sup> C)	25	24	24	25	25
Cloud Cover	Dense Cloud	Overcast	Overcast	Dense Cloud	Dense Cloud
Max RH I (%)	95	95	95	95	95
Min RH II (%)	65	65	65	65	65
Wind speed (km/h)	010	012	008	012	008
Wind direction	SE	SE	SE	SE	SE

Light to moderate rain may likely to occur from 26<sup>th</sup> to 30<sup>th</sup> August 2017 with dense cloudy to overcast sky.

According to Meteorological observations recorded at Agrometeorological Observatory, AMFU-Pantnagar, G B Pant University of Agriculture & Technology (Altitude: 243.8 m) during last seven days (18 – 24 August, 2017) are as follows:

Weather Parameter	Udham Singh Nagar						
	18/08/2017	19/08/2017	20/08/2017	21/08/2017	22/08/2017	23/08/2017	24/08/2017
Rainfall (mm)	19.4	7.0	Trace	0.0	4.2	53.2	0.0
Max Temp ( <sup>0</sup> C)	33.6	33.5	32.6	33.5	32.0	32.6	27.5
Min Temp ( <sup>0</sup> C)	25.5	25.1	26.1	26.5	26.0	26.1	24.0
CC I CC II	Overcast	Overcast	Dense Cloud	Dense Cloud	Dense Cloud	Overcast	Overcast
	Dense Cloud	Cloudy	Partly Cloudy	Cloudy	Cloudy	Overcast	Cloudy
Max RH I (%)	90	93	80	83	86	92	96
Min RH II (%)	74	73	70	73	74	95	74
Wind speed (km/h)	7.0	6.7	7.1	6.8	5.4	5.0	3.3
Wind direction I	ENE	ENE	ENE	ESE	ENE	ENE	N
Wind Direction II	ENE	ENE	ENE	WNW	ESE	ESE	WNW

## Weather Based Agro-Advisories

### Crop Management:

- If the leaves of paddy crops from tip to bottom are drying after turning yellow, then 15 g streptomycin + 500 g Copper Oxochloride in 500 liter of water per hectare should be sprayed.
- On the occurrence of red coloured oval spot on the leaves of maize, solution of 1.5 kg mancozeb per hectare should be sprayed.
- On the occurrence of thin and long light yellow coloured spot which turn kathai later, then solution of 5 kg zinc sulfate + 20 kg urea in 1000 liter of water per hectare should be sprayed.
- Attack of Hispa insect in paddy are also observed in many places. These Hispa insects are black in colour and thorns are found in their body. They consume chlorophyll of the leaves and turn them white. In case of occurrence of Black Hispa insect in paddy, spray 40 EC Triazophos @ 750ml/Ha or 36 SL Monocrotophos @ 1400ml/Ha should be done in the field.
- On the occurrence of Khaira disease in paddy crop, spray 5 Kg Zinc Sulphate and 20 Kg Urea or 2.5 Kg slacked lime, after mixing with 1000 liter of water.
- To control the shoot borer and top borer in Sugarcane, apply Carbofuron 3G @ 30Kg/Ha in the field when sufficient moisture is available in the soil.
- In monsoon season, it is advised that necessary arrangements should be done for drainage of water in the field of Sugarcane crop.

### Horticultural Management:

- If seedlings of Brinjal are ready in the nursery, then transplant them in this month. For transplanting row to row and plant to plant distance should be 60 cm.
- Transplanting of Chilly should be completed in the month of August and it should be transplanted on the ridge. Fast growing varieties of Chilly should be transplanted at a distance of 50\*50 cm and slow growing varieties at a distance of 45\*30 cm.
- Mid variety of Cabbage can be sown in this month and weeding and hoeing should be done in the early variety of Cabbage.
- Field should be prepared for early variety of Spinach (Palak) and sowing should be done in this month.
- Spinach should be sown at 25-30 cm distance in the lines and seed rate should be @25-30kg seeds/Ha.
- In this month, there is a chance of the occurrence of tent caterpillars. So tent should be removed with the help of tent removable device and effected part should be removed and burnt together with insects. If the occurrence of this insect is high then 0.2 % carbaryl or 0.05 % quinalphos should be sprayed.
- To control the red rust and anthracnose, 0.3 % Copper oxi chloride (3.0 g/liter) should be sprayed.
- Establishment of new orchard should be done.

### Animal Husbandry:

- Green fodder should be given in less quantity to the animals. It is advised that green fodder can be given to the animals by mixing it in dry fodder.
- Drinking water should be clean because there is a possibility of occurrence of viruses due to parasite and fungus.
- Animal shed should be dried, for this, lime should be sprayed time to time in their places.
- Deworming doses should be given to the poultry birds on the recommendation of veterinarian because worms in the poultry birds lessen the production capacity of eggs.
- To save the poultry farm from fly and mosquito, Melathian or Finit should be sprayed.

- After animal's delivery, Gynotone or Utrotone medicine @ 200ml should be given in morning and evening hours for three days to clean the womb.

**Dr. R K Singh**  
**Professor & Principal Nodal Officer-GKMS**  
**AMFU- Pantnagar**