

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Nainital

Year: 27 Issue No.: 50 Bulletin Period: 27 June – 1 July 2018 Day: Tuesday Dated: 26.06.2018

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Nainital district are as follows:

Parameter	Nainital				
	27/06/2018	28/06/2018	29/06/2018	30/06/2018	01/07/2018
Rainfall (mm)	10	25	45	60	70
Max Temp ($^{\circ}$ C)	24	23	22	22	21
Min Temp ($^{\circ}$ C)	14	14	13	12	13
Cloud Cover	Dense Cloudy	Dense Cloudy	Dense Cloudy	Overcast	Overcast
Max RH I (%)	80	85	90	95	95
Min RH II (%)	45	50	55	55	60
Wind speed (km/h)	004	006	008	008	006
Wind direction	SE	SE	SE	ESE	SE

Light rain on 27th June and moderate to heavy rain from 28th June to 01st July with dense cloudy to overcast sky may likely to occur.

According to the observations recorded at India Meteorological Department Observatory located at Nainital (Altitude: 2084 m) during last seven days (19-25 June, 2018), it is observed that state of sky were p-cloudy to dense cloudy. Maximum and minimum temperature varied between 24.2 $^{\circ}$ C to 26.0 $^{\circ}$ C and 14.3 $^{\circ}$ C to 17.3 $^{\circ}$ C respectively.

Weather Based Agro-Advisories

Crop Management:

- Nursery of basmati paddy should be prepared. Seed rate for nursery should be 15-20 kg/ha.
- Certified seeds or self-produced healthy seeds should be used for nursery of paddy only after treatment.
- To control the weed in the nursery, solution of 1.5-2 kg Pretilachlor 50EC in 750 lit of water should be sprayed within 3 days. If Pretilachlor is not available then 1.3 lit Anilofos 30 EC or 200-250ml Bispyribac Sodium 10 EC (Nominee gold) should be used after 10-15 days of sowing.
- Transplanting of nursery, sown earlier, should be done till 15th July.
- If the leaves of french bean and lobia are drying, carbendazim @ 1 g/lit of water should be used.


Horticultural Management:

- For grafting of temperate fruits, T-budding or chip budding process should be started.

- In mid hills, plucking the early varieties of apple should be started and send them to the market.
- To control the gummosis disease in fruit bearing fruit trees, Streptocycline 0.01% or copper oxichloride 0.025% should be sprayed at 15 days interval.
- In mid hilly regions, to save the late maturing fruit varieties of apple from falling, plenofix @ 10ppm should be sprayed.
- In high hilly regions, to control the scab disease on the falling of pankhuri of flower in apple crop, Carbendazim 0.05 % should be used.
- In mid hilly regions, short duration variety of cabbage, cauliflower, broccoli and knol-khol should be selected and sown in the nursery.
- If proper moisture is available in the fields then sowing of radish, rai, coriander, carrot etc should be done.
- In mid hilly regions, weeding, hoeing and irrigation should be done in the crops sown in the polyhouse and recommended chemical should be sprayed to save the crop from insects.
- In the standing crop of cucumber and chappan kaddu necessary irrigation should be given after weeding and hoeing in mid hilly regions. Ready fruits should be sent to the market.
- In valley, digging of potato, onion and garlic should be done if crops are ready.
- In valley, picking up the green pods of French bean should be done.
- In valley, to control the insects in the standing crop of tomato and capsicum, Melathiyam (15ml/10liter of water) and to control the blight disease, Mancozeb (20g/10liter of water) should be sprayed in the day time under proper sunny condition.
- In mid hilly regions, to control the blight disease in potato crop, Mancozeb should be sprayed.
- To control the leaf blight disease in onion, tebuconazole or difinoconazole or propiconazole @ 500ml/ha with any sarvangi insecticide and sticker should be sprayed.
- On the occurrence of contracted piebald leaves in chilly and tomato, infected plants should be removed and destroyed. To control the juice sucking insect in these crops, sarvangi insecticide should be sprayed. To save the crop from late blight disease, it is advised that solution of Mancozeb @ 2.5g/liter or copper oxichloride @ 3g/lit of water should be sprayed.

Animal Husbandry:

- To save the animals from infected disease, vaccination should be done on the recommendation of veterinarian.
- On the occurrence of nitrate toxicity in the food of animals, respiration and pulse rate of animals increases and they feel weakness and stiffness in their muscles. Give the 50-100 ml dose of 1% Methylene blue direct in the veins of animals if their food is affected by nitrogen toxicity.
- If animal has consumed the cyanide affected fodder then water should not be given to them. Animals should be kept away from grazing the less grown jawar, bajara and fodder in pasture.
- Small shrivel/fade yellow stiff plants should not be used as fodder for animals to save them from cyanide.
- Neat, fresh and cold water should be given to the animals three times in a day. Availability of sufficient moisture in the body of animals maintains the balance between the temperature of their skin and environment which save the animals from the effect of loo (summer wind).



Dr. R K Singh

**Professor & Principal Nodal Officer-GKMS
AMFU- Pantnagar**