

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Udham Singh Nagar

Year: 26 Issue No.: 75 Bulletin Period: 27 Sept – 1 Oct, 2017 Day: Tuesday Dated: 26-09-2017

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Udham Singh Nagar and plain area of Nainital districts are as follows:

Parameter	Udham Singh Nagar				
	27/09/2017	28/09/2017	29/09/2017	30/09/2017	01/10/2017
Rainfall (mm)	0	0	0	0	0
Max Temp (⁰ C)	33	34	34	34	35
Min Temp (⁰ C)	23	24	24	24	24
Cloud Cover	Cloudy	P-Cloudy	P-Cloudy	P-Cloudy	P-Cloudy
Max RH I (%)	90	90	85	85	85
Min RH II (%)	60	60	55	55	55
Wind speed (km/h)	006	006	006	008	008
Wind direction	NW	ESE	ESE	ESE	ESE

According to Meteorological observations recorded at Agrometeorological Observatory, AMFU-Pantnagar, G B Pant University of Agriculture & Technology (Altitude: 243.8 m) during last seven days (19 - 25 Sept, 2017) are as follows:

Weather Parameter	Udham Singh Nagar						
	19/08/2017	20/08/2017	21/08/2017	22/09/2017	23/09/2017	24/09/2017	25/09/2017
Rainfall (mm)	76.2	0.0	0.0	26.6	91.2	7.2	0.0
Max Temp (⁰ C)	35.0	32.5	34.9	31.3	25.2	29.0	31.5
Min Temp (⁰ C)	22.0	25.0	24.4	24.9	23.3	24.0	23.5
CC I	Overcast	Clear	Dense Cloud	Overcast	Dense Cloud	Overcast	Dense Cloud
CC II	Cloudy	Cloudy	Cloudy	P-Cloudy	P-Cloudy	Overcast	Cloudy
Max RH I (%)	95	87	84	95	95	93	95
Min RH II (%)	71	62	78	93	86	77	72
Wind speed (km/h)	2.9	4.9	3.6	4.8	7.3	9.5	5.7
Wind direction I	ENE	Calm	ENE	ESE	ENE	ENE	WNW
Wind Direction II	WSW	W	ESE	ESE	WSW	WSW	WSW

Weather Based Agro-Advisories

Crop Management:

- To control the bacterial leaf blight disease in paddy crop, streptomycin 15 g + copper oxichloride 500 g / hectare should be sprayed.
- If growth of Sugarcane is good then tie the plants of three basins of the two rows by using Kaichi method.
- On the occurrence of yellow mosaic disease in the Pulse (Dalhania) crops, leaves turn yellow. This disease is spread by the White Fly. To control this, 2-3 spraying of 30 EC Dimethoate or 20 EC Methyl-O-dimeton @ 1liter/Ha in 500-600 liter of water should be done at an interval of 10-12 days.
- In no rainfall situation, irrigation should be given at the time of flowering in Pulse crops and if there is shortage of moisture in the Cowpea then one irrigation must be applied at the time of fruiting.
- On the attack of stem borer, Chlorantraniliprole 20 SC @ 150ml/ha or Kartop 50SP @ 1kg/Ha or Flubendiamide 480 SC @75ml/Ha or Monocrotophos 36 SL @ 1400 ml/Ha should be sprayed.
- Attack of Hispa insect in paddy are also observed in many places. These Hispa insects are black in colour and thorns are found in their body. They consume chlorophyll of the leaves and turn them white. In case of occurrence of Black Hispa insect in paddy, spray 40 EC Triazophos @ 750ml/Ha or 36 SL Monocrotophos @ 1400 ml/Ha should be done in the field.
- To control the shoot borer and top borer in Sugarcane, apply Carbofuron 3G @ 30Kg/Ha in the field when sufficient moisture is available in the soil.

Horticultural Management:

- On the occurrence of leaf spot disease in brinjal crop, solution of mancozeb @ 2g/lit of water should be sprayed.
- To control the sucking insect in the Colocasia, 0.2 % solution of Imidachlorpid + Dithene 45 should be sprayed on 10 days interval.
- Weeding and hoeing should be done in the Brinjal crop, sown in the previous month. 1/4th part of Nitrogen should be applied as top dressing in the field and remaining 1/4th part should be given at an interval of 60-65 days as a top dressing in the standing crop.
- To save the Cauliflower from attack of insect, 0.1 % Metasystox or Imidachlorpid should be sprayed.
- After withdrawal of monsoon season, it is observed that Gum comes out from the Mango tree and this is mostly occurred in the orchards where soil is sandy. For this, Copper Sulphate @ 200-250g /Tree should be used. Remove the Gum from the trees by scratching, then spraying of Copper-Oxichloride @ 2g/lit should be done on that part of the tree.
- On the occurrence of branch knot in the Orchard, Quinalphos @ 2ml/liter should be used.
- Clearing and ploughing of orchard should be done.
- For proper fruiting in Mango, use Paclobutrazol. Apply 1ml Active Ingradient (AI) /meter of canopy.
- Continue the transplanting of plant in the orchards.

Animal Husbandry:

- Just after delivery, tie the navel of new born baby of cattle by cutting it with sharp and clean knife, then apply Gentian violet paint or tincture iodine.

- In green fodder, leguminous fodder is best diet for animals and to increase production. So it is advised that cattleman should grow leguminous fodder (berseem) to keep their cattle healthy and to obtain maximum production from them.
- In this month, rate of animal's delivery especially in buffalos increases. The Pregnant animals should be separated from other animals if possible and complete diet in little quantity at many times in a day should be given to them otherwise they can be affected by afra (fatty liver).
- Within 2 hours of animal's delivery, little quantity of colostrums should be given to the new born baby of cattle after their proper cleaning.
- Green fodder should be given in less quantity to the animals. It is advised that green fodder can be given to the animals by mixing it in dry fodder.
- Drinking water should be clean because there is a possibility of occurrence of viruses due to parasite and fungus.
- Animal shed should be dried, for this, lime should be sprayed time to time in their places.
- Deworming doses should be given to the poultry birds on the recommendation of veterinarian because worms in the poultry birds lessen the production capacity of eggs.

Dr. R K Singh
Professor & Principal Nodal Officer-GKMS
AMFU- Pantnagar