

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



## Gramin Krishi Mausam Sewa Bulletin, District - Nainital

Year: 27 Issue No.: 59 Bulletin Period: 28 July – 1 Aug, 2018 Day: Friday Dated: 27.07.2018

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Nainital district are as follows:

Parameter	Nainital				
	28/07/2018	29/07/2018	30/07/2018	31/07/2018	01/08/2018
Rainfall (mm)	25	25	40	60	40
Max Temp ( $^{\circ}$ C)	18	19	20	21	22
Min Temp ( $^{\circ}$ C)	15	14	13	13	14
Cloud Cover	Dense Cloudy				
Max RH I (%)	95	95	95	90	90
Min RH II (%)	50	50	55	50	50
Wind speed (km/h)	006	008	006	004	004
Wind direction	ENE	ESE	ENE	SE	E

**Moderate to heavy rain with dense cloudy sky may likely to occur in coming five days.**

According to the observations recorded at India Meteorological Department Observatory located at Nainital (Altitude: 2084 m) during last seven days (20 - 26 July, 2018), it is observed that state of sky were cloudy to dense cloudy. Maximum and minimum temperature varied between  $18.5^{\circ}$ C to  $23.6^{\circ}$ C and  $14.9^{\circ}$ C to  $15.8^{\circ}$ C respectively.

### Weather Based Agro-Advisories

#### Crop Management:

- Transplanting of paddy should be completed in this month.
- Only 2-3 seedlings should be transplanted at one place.
- Line to line and plant to plant distance for transplanting paddy should be 20-25 cm and 10cm respectively.
- To control the weed in transplanted paddy, solution of pretilachlor 50 EC @ 1.5-2.0 lit/500 lit of water or solution of butachlor (50 EC) @ 3 liter/ 750 liter of water should be sprayed within 3 days of transplanting. *or*
- Solution of cyhalofop-butyl 10 EC @ 1lit/500 lit of water should be used to control the weed in transplanted paddy, after 20-25 days of transplanting.

- Bispyribac Sodium 10EC (Nominee gold) @ 200-250 ml /500 liter of water should be sprayed after 20-25 days of transplanting. *or* Metsulfuron-methyl 10% + chlorimuron ethyl 10% WP @ 20g/500liter of water should be sprayed after 20-25 days of transplanting or on the emergence of 3-4 leaves. It controls the broad leaf and motha class weeds.
- Weeds should be removed from the sugarcane field and irrigation and drainage should be maintained.


### **Horticultural Management:**

- To control the root and stem joint rot disease in tomato and chilly, irrigate the roots of the crops with the solution of trichoderma @ 10 g/lit or carbendazim @1 g/lit of water.
- Radish, spinach, rai, turnip should be sown.
- Cucumber and chappan kaddu should be sent to the market.
- If the upper stipe of chilly crop is drying after turning black, then infected branches should be removed by plucking to save the crop and to save the crop from rotting solution of 0.1% Carbendazim should be sprayed.
- On the occurrence of spot on the leaves of tomato and scorching of leaves, solution of mancozeb @ 2.5 g/lit of water should be sprayed.
- On the rotting of capsicum, solution of mancozeb @ 2.5 g/lit of water should be sprayed.
- If growth of white fungus is observed and there is the rotting of pods in French bean, solution of carbendazim @ 1g/lit of water should be sprayed.
- In hilly regions, if moisture in the soil is more than sufficient, then sowing of radish, rai, coriander, turnip and spinach (Pusa Harit) may be done.
- In hilly regions, tomato, Brinjal and Capsicum crop, proper drainage should be maintained in the field during monsoon season and fruit picking should be done timely.
- In case of heavy rainfall, it is advised that proper drainage should be made from the basins of fruit plant.
- Evergreen fruit plants like Mango, guava, Lemon, papaya, Litchi etc should be planted in this month.
- Weeding and hoeing operation may be taken up in the Orchard.
- To control the gummosis disease in fruit bearing fruit trees, Streptocycline 0.01% or copper oxichloride 0.025% should be sprayed at 15 days interval.
- In mid hilly regions, to save the late maturing fruit varieties of apple from falling, plenofix @ 10ppm should be sprayed.
- Grafting of mango, guava, lemon and litchi fruits should be started.
- Drainage of water should be done in the basins of apple.

### **Animal Husbandry:**

- After animal's delivery, Gynotone or Utrotone medicine @ 200ml should be given in morning and evening hours for three days to clean the womb.
- Rain water should not be given to the animals.
- To save the animals from infected disease, vaccination should be done before onset of monsoon on the recommendation of veterinarian.
- July month is highly favourable for cattles' delivery. So, it is recommended that their dwelling place should be clean and hygienic.
- Just after delivery, tie the navel of new born baby of cattle by cutting it with sharp and clean knife, then apply Gentian violet paint or tincture iodine.

- Don't leave the animal in the monsoonal rain because this may lead the irritation in the skin of animals.
- Make the necessary arrangements to save the animals during adverse weather.
- Give Vaccination to save the animals from infectious diseases before onset of monsoon.
- Give deworming to the animals on the recommendation of veterinarian.
- Don't give the high dose of green fodder to the horses because high dose of this increase the chances occurrence of Celiac disease.

A handwritten signature in black ink, appearing to read 'R K Singh', with a horizontal line underneath.

**Dr. R K Singh**  
**Professor & Principal Nodal Officer-GKMS**  
**AMFU- Pantnagar**