

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Udham Singh Nagar

Year: 27 Issue No.: 94 Bulletin Period: 28 Nov – 2 Dec, 2018 Day: Tuesday Dated: 27-11-2018

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Udham Singh Nagar district are as follows:

Parameter	Udham Singh Nagar				
	28/11/2018	29/11/2018	30/11/2018	01/12/2018	02/12/2018
Rainfall (mm)	0	0	0	0	0
Max Temp (⁰ C)	27	26	26	25	25
Min Temp (⁰ C)	11	10	9	8	8
Cloud Cover	Cloudy	Cloudy	P-Cloudy	Clear	Clear
Max RH I (%)	85	90	85	90	90
Min RH II (%)	45	45	40	45	45
Wind speed (km/h)	006	004	006	006	006
Wind direction	ESE	ESE	NW	NW	NW

According to Meteorological observations recorded at Agrometeorological Observatory, AMFU-Pantnagar, G B Pant University of Agriculture & Technology (Altitude: 243.8 m) during last seven days (20 – 26 Nov, 2018) are as follows:

Weather Parameter	Udham Singh Nagar						
	20/11/2018	21/11/2018	22/11/2018	23/11/2018	24/11/2018	25/11/2018	26/11/2018
Rainfall (mm)	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Max Temp (⁰ C)	25.6	27.5	27.0	26.5	26.0	25.5	28.0
Min Temp (⁰ C)	10.9	9.9	11.4	9.4	8.9	10.8	10.4
CC I	0	0	0	0	0	0	0
CC II	0	0	0	0	0	0	0
Max RH I (%)	95	90	93	95	95	93	75
Min RH II (%)	50	57	46	46	46	57	57
Wind speed (km/h)	1.2	1.9	1.8	2.1	2.3	1.7	1.1
Wind direction I	Calm	Calm	Calm	Calm	Calm	N	Calm
Wind Direction II	WNW	WNW	WNW	WSW	WNW	WSW	WNW

Weather Based Agro-Advisories

Crop Management:

- In timely sown wheat, first irrigation should be given after 20-25 days of sowing. After 3-4 days of first irrigation, remaining dose of Nitrogen should be applied as top dressing in the afternoon.
- To control the weed in the wheat, first and second weeding and hoeing should be done after 25-30 days and 45-50 days of sowing respectively.
- In irrigated condition, late sowing of gram should be completed in the first week of December. Pusa-372 and PG 186 should be selected and sow at the depth of 6-8cm in the lines at a distance of 20-25cm.
- Weeding and hoeing is necessary two times in the timely sown gram crop. First weeding and hoeing should be done after 25-30 days of sowing and second 45-50 days after sowing and after first irrigation.
- Sowing of lentil should be completed up to 15th of December and weeding and hoeing should be done after 25-30 days of sowing.
- Harvesting of Arhar should be done.
- Check the quantity of brix in the juice of ratoon sugarcane crop. When quantity of brix reaches up to 18%, then cutting of sugarcane should be done and send them to the mill. Immediately after the harvesting of sugarcane, field should be prepared for rabi crops.


Horticultural Management:

- Radish can be sown throughout the year. Seed rate for European and Asian variety of Radish should be 6-8Kg/ha and 10-12 kg/Ha respectively.
- Dung manure should be applied in the field before sowing the seed of radish. Fertilizer should be used on the basis of soil testing and distance of line of sowing should be 20-25cm. On the germination of seed plant to plant distance should be maintained at a distance of 8-10 cm.
- Weeding and hoeing should be done in the garlic crop.
- If the seedlings of late variety of cauliflower are ready then its transplanting should be done in this month. 150 kg of N, 80 kg of P and 60 kg of K are recommended for transplanting, from which half dose of nitrogen, full dose of phosphorous & potash should be used at the time of final ploughing of the field.
- If gum are releasing from the stem of the tree then copper oxichloride and slaked lime should be apply in the infected areas of main stem.
- On the occurrence of anthracnose on the leaves of mango, copper oxichloride @ 2g/liter should be sprayed.
- On the occurrence of web insect, cleaning of web should be done by using web machine and Quinalphos @ 2ml/liter should be sprayed.
- If stem canker or kai kawak are sown in the stem of mango tree then copper oxichloride or beedo mixer should be sprayed.
- Clearing and ploughing of orchard should be done at the earliest.

Animal Husbandry:

- Vaccination of FMD (Foot and Mouth Disease) should be given to the animals.
- Foot and Mouth Disease is identified by red eyes, high fever, less productivity and food taken capacity, Mouth ulcers, wound in foot due to delay in the treatment of animals. Infected animals should be separated from healthy animals.

- Keeping in mind the probable high rate of Cattle's delivery in this month, especially in Buffalo, animal shed should be well neat, clean, dry and ventilated. For this, dry Lime should be sprayed in the nearby places.
- In case of new born, placenta should be cut with new blade and tie a knot in that. Don't forget to apply Betadine or Tincher on the cut.
- Toxocara vitulorum (patera) are found in the food tube of buffalo calf in the age of 1-4month. More than 50% buffalo calf should be died because of delay in treatment. Piperazine should be given to the infected calf on identification of Patera disease.
- To save the buffalo calf from patera disease, 10-15 CC Neem oil should be given to the calf immediately after 10 days of his birth. Again after 10 days, 10-15 CC Neem oil should be given. Chenopodium album oil can also be given as a fodder to animals to save the calf/animals from patera disease.
- It is advised that green fodder can be given to the animals by mixing it in dry fodder otherwise animal may be infected by Timpeti disease which causes death of animals.
- There is a chance of Aflatoxicosis disease in Poultry due to occurrence of fungus in the food which causes to death of poultry birds. So, medicine should be administered to the poultry birds only on the advice of veterinarian.



Dr. R K Singh
Professor & Principal Nodal Officer-GKMS
AMFU- Pantnagar