

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Nainital

Year: 27 Issue No.: 68 Bulletin Period: 29 Aug – 2 Sept, 2018 Day: Tuesday Dated: 28.08.2018

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Nainital district are as follows:

Parameter	Nainital				
	29/08/2018	30/08/2018	31/08/2018	01/09/2018	02/09/2018
Rainfall (mm)	25	20	15	35	25
Max Temp ($^{\circ}$ C)	20	21	22	21	20
Min Temp ($^{\circ}$ C)	15	14	15	14	15
Cloud Cover	Dense Cloudy				
Max RH I (%)	85	85	85	90	95
Min RH II (%)	45	45	45	50	50
Wind speed (km/h)	006	002	004	006	004
Wind direction	SE	ESE	SE	E	ESE

Light to moderate rain with dense cloudy sky may likely to occur in coming five days.

According to the observations recorded at India Meteorological Department Observatory located at Nainital (Altitude: 2084 m) during last seven days (21 – 27 Aug, 2018), it is observed that state of sky were cloudy to dense cloudy. Maximum and minimum temperature varied between 16.7° C to 21.2° C and 15.0° C to 16.0° C respectively.

Weather Based Agro-Advisories

Crop Management:


- To control the leaf blight disease in maize crop, mancozeb @ 2.5g/lit of water should be sprayed.
- Remaining dose of nitrogen should be given as a top dressing twice at the time of panicle initiation after 20 days of transplanting and starting stage of ear formation i.e. 40-50 days of transplanting. Proper moisture should be available in the field at the time of top dressing and rain drops are not there in the leaves.
- Arrangement of proper drainage should be done in pulses and sugarcane crops.
- Weeds should be removed from the sugarcane field and irrigation and drainage should be maintained.

Horticultural Management:

- In valley, cabbage and cauliflower snowball should be transplanted in irrigated condition for domestic consumption.
- In high hills, sowing of Arkil variety of pea should be done.
- To control the white rotting in french bean and cowpea (lobia), Carbendazim 1 g/liter should be sprayed.
- To control the root and stem joint rot disease in tomato and chilly, irrigate the roots of the crops with the solution of trichoderma @ 10 g/lit or carbendazim @1 g/lit of water.
- If the upper stipe of chilly crop is drying after turning black, then infected branches should be removed by plucking to save the crop and to save the crop from rotting solution of 0.1% Carbendazim should be sprayed.
- On the occurrence of spot on the leaves of tomato and scorching of leaves, solution of mancozeb @ 2.5 g/lit of water should be sprayed.
- On the rotting of capsicum, solution of mancozeb @ 2.5 g/lit of water should be sprayed.
- If growth of white fungus is observed and there is the rotting of pods in French bean, solution of carbendazim @ 1g/lit of water should be sprayed.
- In case of heavy rainfall, it is advised that proper drainage should be made from the basins of fruit plant.
- Evergreen fruit plants like Mango, guava, Lemon, papaya, Litchi etc should be planted in this month.
- Weeding and hoeing operation may be taken up in the Orchard.
- To control the gummosis disease in fruit bearing fruit trees, Streptocycline 0.01% or copper oxichloride 0.025% should be sprayed at 15 days interval.

Animal Husbandry:

- Green fodder should be given in less quantity to the animals. It is advised that green fodder can be given to the animals by mixing it in dry fodder.
- Drinking water should be clean because there is a possibility of occurrence of viruses due to parasite and fungus.
- Animal shed should be dried, for this, lime should be sprayed time to time in their places.
- Deworming doses should be given to the poultry birds on the recommendation of veterinarian because worms in the poultry birds lessen the production capacity of eggs.
- To save the poultry farm from fly and mosquito, Melathiyan or Finit should be sprayed.
- After animal's delivery, Gynotone or Utrotone medicine @ 200ml should be given in morning and evening hours for three days to clean the womb.



Dr. R K Singh

**Professor & Principal Nodal Officer-GKMS
AMFU- Pantnagar**