

ग्रामोभ्युदयादेव देशोभ्युदयः

Govind Ballabh Pant University of Agriculture and Technology

Department of Agrometeorology

College of Agriculture

Pantnagar-263145, Uttarakhand

Phone No. 05944 - 233 032



Gramin Krishi Mausam Sewa Bulletin, District - Nainital

Year: 27 Issue No.: 77 Bulletin Period: 29 Sept – 3 Oct, 2018 Day: Friday Dated: 28.09.2018

As per forecast received from National Centre for Medium Range Weather Forecast, India Meteorological Department, Mausam Bhawan, New Delhi and value added to the forecast by Meteorological Centre, Dehradun for next five days for Nainital district are as follows:

Parameter	Nainital				
	29/09/2018	30/09/2018	01/10/2018	02/10/2018	03/10/2018
Rainfall (mm)	1	0	0	0	0
Max Temp ($^{\circ}$ C)	22	22	22	23	23
Min Temp ($^{\circ}$ C)	12	12	12	13	13
Cloud Cover	Cloudy	P-Cloudy	P-Cloudy	P-Cloudy	P-Cloudy
Max RH I (%)	85	80	75	70	70
Min RH II (%)	45	40	35	35	35
Wind speed (km/h)	004	006	006	004	004
Wind direction	NW	NW	NW	NW	NW

Weather will remain clear with p-cloudy sky in coming five days.

According to the observations recorded at India Meteorological Department Observatory located at Nainital (Altitude: 2084 m) during last seven days (21 – 27 Sept, 2018), it is observed that state of sky were cloudy to dense cloudy. Maximum and minimum temperature varied between 13.8° C to 21.4° C and 10.5° C to 14.0° C respectively.

Weather Based Agro-Advisories

Crop Management:

- Sowing of winter season sugarcane crop should be done up to 15th October.
- Farmers who didn't complete the sowing of toria crop till now, they can complete it up to first week of October.
- For sowing of toria, seed rate and row to row distance should be 5kg/ha and 30cm respectively.
- Seed treatment of toria should be done with Thiram 2.5g/kg or Mancozeb 3g/kg.
- For toria, 90kg Nitrogen, 40 kg phosphorous and 20 kg Potash are used. Full dose of phosphorous, potash and half dose of nitrogen should be used at the time sowing. Remaining half dose of nitrogen should be used after 20-25 days of sowing.


- To control the brown leaf spot in paddy crop, solution of mancozeb @ 2.5 g/lit of water should be sprayed.
- In paddy crop, weed management and proper storage of water should be done.
- To control the leaf blight disease in maize crop, mancozeb @ 2.5g/lit of water should be sprayed.

Horticultural Management:

- In mid and high hills, basin should be used in pea, radish, cabbage, cauliflower crops to conserve the moisture after hoeing.
- In valley, cabbage and cauliflower snowball should be transplanted in irrigated condition for domestic consumption.
- To control the root and stem joint rot disease in tomato and chilly, irrigate the roots of the crops with the solution of trichoderma @ 10 g/lit or carbendazim @1 g/lit of water.
- If growth of white fungus is observed and there is the rotting of pods in French bean, solution of carbendazim @ 1g/lit of water should be sprayed.
- Evergreen fruit plants like Mango, guava, Lemon, papaya, Litchi etc should be planted in this month.
- Weeding and hoeing operation may be taken up in the Orchard.

Animal Husbandry:

- Just after delivery, tie the navel of new born baby of cattle by cutting it with sharp and clean knife, then apply Gention violet paint or tincture iodine.
- In green fodder, leguminous fodder is best diet for animals and to increase production. So it is advised that cattleman should grow leguminous fodder (berseem) to keep their cattle healthy and to obtain maximum production from them.
- In this month, rate of animal's delivery especially in buffalos increases. The Pregnant animals should be separated from other animals if possible and complete diet in little quantity at many times in a day should be given to them otherwise they can be affected by afra (fatty liver).
- Within 2 hours of animal's delivery, little quantity of colostrums should be given to the new born baby of cattle after their proper cleaning.
- Green fodder should be given in less quantity to the animals. It is advised that green fodder can be given to the animals by mixing it in dry fodder.
- Vaccination of FMD (Foot and Mouth Disease) should be given to the animals.



Dr. R K Singh

Professor & Principal Nodal Officer-GKMS

AMFU- Pantnagar