2/7/25, 5:23 PM District Advisory



# Gramin Krishi Mausam Sewa Agromet Advisory Bulletin for NAINITAL District

# Agromet Advisory Bulletin for NAINITAL District Govind Ballabh Pant University of Agriculture & Technology



Udham Singh Nagar, Pantnagar, Uttarakhand

# **Agromet Advisory Bulletin**

Date: 2025-02-07

Weather Forecast of District NAINITAL (Uttarakhand) Issued On: 2025-02-07 (Valid Till 08:30 IST of the next 5 days)

Parameter	2025-02-08	2025-02-09	2025-02-10	2025-02-11	2025-02-12
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	16.0	17.0	18.0	18.0	18.0
Tmin(°C)	4.0	5.0	5.0	6.0	5.0
RH-I(%)	30	35	35	35	40
RH-II(%)	25	30	30	30	25
Wind Speed(kmph)	8	8	7	6	7
Wind Direction(Degree)	320	330	320	310	320
Cloud Cover(Octa)	2	4	2	1	1

#### **Weather Summary/Alert:**

In the upcoming days no rainfall has been predicted. Maximum & minimum temperature may be 16.0-18.0 degree Celsius & 5.0- 6.0 degree Celsius, respectively. Wind is expected to blow with a speed of 6-8 km/hr primarily from the west -north-west and north-west direction. Dry weather is likely to prevail.

#### **General Advisory:**

A regular update regarding weather forecast and agrometeorological advisory is available on the "Meghdoot App" while "Damini App" is also available to get lightning information. Meghdoot and Damini apps can be downloaded from Google Play Store (Android users) and App Center (iOS users). The NDVI composite shows moderate agricultural vigour in the region. The extended range forecast system shows large deficient rainfall and below normal maximum and normal minimum temperature trend during 7 to 13 February.

#### **SMS Advisory:**

As per IMD forecast, dry weather is expected so irrigation should be applied and the sprays against disease-pest should be scheduled accordingly.

#### **Crop Specific Advisory:**

Crop (Stage)	Crop Specific Advisory
LENTIL	The crop should be regularly monitored for weeding and hoeing while irrigation should be applied as per requirement. Occurrence of plant rot can be treated with appropriate fungicide application.
RAPESEED	The crop should be monitored regularly and time to time irrigation should be provided. Diseases should be monitored and control measures should be taken up.
WHEAT	Irrigation should be applied as per requirement and urea application should be done after irrigation.
BARLEY	Urea should be used after rainfall and after getting moisture as per requirement.

### **Horticulture Specific Advisory:**

Horticulture (Stage)	Horticulture Specific Advisory
CAPSICUM	Seeds should be sown in the prepared nursery and seedlings should be protected from frost and extreme cold conditions by using plastic covers with proper air and water movement.
VEGETABLE PEA	The crop should be regularly monitored for weeding and hoeing while irrigation should be applied as per requirement. Occurrence of plant rot can be treated with appropriate fungicide application.
SPINACH	Time to time irrigation to avoid chilling injury should be applied and proper measures to protect them against cold conditions should be taken up.
РОТАТО	To control the late blight disease in potato, it is advised that solution of Mancozeb@ 2.5g/liter of water should be sprayed.
APPLE	To control the fruit Rot disease in apple and seed bearing fruits, soil should be removed from the surrounding area of the stem so that the sunrays may enter directly in the stem of the affected tree. Affected bark should be removed and chaubatia paste should be applied along with soil covering.

# **Live Stock Specific Advisory:**

Live Stock	Live Stock Specific Advisory
COW	To save the animal from cold, increase the amount of oil and Jaggery (Gur) in the food of animals. Caraway (Ajwain) and gur also be given to the animals. Vaccination should be given to the animals to save them from Rinderpest disease (Sheetla rog). The animal feed should be increased by 10% for extra energy requirement.
BUFFALO	To save the animal from cold, increase the amount of oil and Jaggery (Gur) in the food of animals. Caraway (Ajwain) and gur also be given to the animals. Vaccination should be given to the animals to save them from Rinderpest disease (Sheetla rog). The animal feed should be increased by 10% for extra energy requirement.

# **Poultry Specific Advisory:**

Poultry	Poultry Specific Advisory		
HHN	Medicine as per the veterinarian should be administered for aflatoxiosis caused due to fungal growth in their feed.		