Gramin Krishi Mausam Sewa Agromet Advisory Bulletin for UDAMSINGH-NAGAR District Govind Ballabh Pant University of Agriculture & Technology Udham Singh Nagar,Pantnagar, Uttarakhand



Agromet Advisory Bulletin

Date : 2024-11-08

Weather Forecast of District UDAMSINGH-NAGAR (Uttarakhand) Issued On : 2024-11-08 (Valid Till 08:30 IST of the next 5 days)

Parameter	2024-11-09	2024-11-10	2024-11-11	2024-11-12	2024-11-13
Rainfall(mm)	0.0	0.0	0.0	0.0	0.0
Tmax(°C)	30.0	30.0	29.0	29.0	29.0
Tmin(°C)	17.0	17.0	16.0	16.0	16.0
RH-I(%)	70	70	70	70	70
RH-II(%)	30	30	30	30	30
Wind Speed(kmph)	2	3	2	2	3
Wind Direction(Degree)	180	160	200	230	230
Cloud Cover(Octa)	1	2	2	1	2

Weather Summary/Alert:

Last week (1st to 7th November) the region did not receive any rainfall and maximum-minimum temperatures ranged from 29.0 to 31.5oC and 16.9 to 21.3oC respectively. The morning and evening relative humidity ranged between 83-94% and 49-62% respectively while the wind was blowing from south-south-east, north-north-west, north-west, west-north-west, north and east-north-east at a speed of 0.9-2.3 kmph. The weather was clear on most of the days last week. As per the upcoming forecast, there will be no rainfall between 9-12th November and maximum-minimum temperatures are expected to be 29.0-30.0oC and 16.0-17.0oC respectively. The wind is expected to blow from south, south-west at a speed of 2-3 kmph. Dry weather likely to prevail from 08-12th November.

General Advisory:

For regular updates on weather conditions in the region, the farmers can get updates from the app "Meghdoot" and lightning update from the app "Damini" available on Google Play store (Android users) and App Centre (iOS users). The NDVI shows good agricultural vigour in the isolated regions of the state ranging 0.35-0.60. The extended range forecast shows deficient rainfall and normal maximum and normal minimum temperature trend during 08.11.2024 to 14.11.2024.

SMS Advisory:

As per IMD forecast, dry weather is expected in the upcoming week so harvested produce should be dried and stored well while sowing operations should be carried in rabi crops.

Crop Specific Advisory:

Crop (Stage)	Crop Specific Advisory
	On ripening the crop should be dug out and after proper drying should be stored well.
PIGEON PEA (RED GRAM/ARHAR)	The pest management measures should be taken up in the crop and if 75-80% pods have ripened in early varieties then the crop should be harvested.

District Advisory

Crop (Stage)	Crop Specific Advisory
CHICK PEA	In last month sown crop, under rainfed conditions should be regularly monitored and weeding should be done at 25-30 days old crop while under irrigated conditions the crop sowing should be done till first fortnight of the month.
LENTIL	In last month sown crop, under rainfed conditions should be regularly monitored and weeding should be done at 25-30 days old crop while under irrigated conditions the crop sowing should be done till first fortnight of the month.
RAPESEED	Regular monitoring of the timely sown crop is important and should be irrigated as and when required before flowering while sowing should be completed.
MUSTARD	Regular monitoring of the timely sown crop is important and should be irrigated as and when required before flowering while sowing should be completed.
WHEAT	Under rainfed conditions the crop should be sown till first week of the month and under irrigated conditions till first fortnight of the month.
FIELD PEA	In last month sown crop, under rainfed conditions should be regularly monitored and weeding should be done at 25-30 days old crop while under irrigated conditions the crop sowing should be done till first fortnight of the month.

Horticulture Specific Advisory:

Horticulture (Stage)	Horticulture Specific Advisory	
ONION	The sowing of rabi onion crop can be done in this month.	
	Sowing of potato can be taken up in this month such as Kufri Badshah, Kufri Jyoti, Kufri Alankar, Kufri Chandranukhi, etc	

Live Stock Specific Advisory:

Live Stock	Live Stock Specific Advisory
BUFFALO	Vaccination of FMD (Foot and Mouth Disease) should be given to the animals. Foot and Mouth Disease is identified by red eyes, high fever, less productivity and food taken capacity, Mouth ulcers, wound in foot due to delay in the treatment of animals. Infected animals should be separated from healthy animals.
COW	Vaccination of FMD (Foot and Mouth Disease) should be given to the animals. Foot and Mouth Disease is identified by red eyes, high fever, less productivity and food taken capacity, Mouth ulcers, wound in foot due to delay in the treatment of animals. Infected animals should be separated from healthy animals.

Others (Soil / Land Preparation) Specific Advisory:

Others (Soil / Land Preparation)	Others (Soil / Land Preparation) Specific Advisory
	The standing late sown kharif crops such as rice, urd, green gram, soyabean and till should be harvested so that rabi crops can be timely sown.