



**Gramin Krishi Mausam Sewa**  
**District Level Agromet Advisory Bulletin**  
**Govind Ballabh Pant University of Agriculture & Technology**



Udham Singh Nagar, Pantnagar, Uttarakhand

## Agromet Advisory Bulletin

Date : 27-08-2021

**Weather Forecast of UDAMSINGH-NAGAR(Uttarakhand) Issued On : 2021-08-27(Valid Till 08:30 IST of the next 5 days)**

Parameter	2021-08-28	2021-08-29	2021-08-30	2021-08-31	2021-09-01
<b>Rainfall</b>	15.0	10.0	8.0	5.0	3.0
<b>Tmax(°C)</b>	30.0	29.0	30.0	31.0	32.0
<b>Tmin(°C)</b>	25.0	24.0	24.0	25.0	25.0
<b>RH-I(%)</b>	90	90	95	90	85
<b>RH-II(%)</b>	65	60	60	60	55
<b>Wind Speed(kmph)</b>	10.0	12.0	12.0	12.0	12.0
<b>Wind Direction(Degree)</b>	100	120	120	140	140
<b>Cloud Cover(Octa)</b>	8	8	5	3	6

### Weather Summary/Alert:

In the last seven days (20 – 26 August, 2021), there was 60.4 mm rain and the sky was cloudy to dense cloudy. The maximum & minimum temperatures ranged between 29.0 to 34.0°C and 24.4 to 26.3°C respectively. Relative humidity ranges from 87 to 97 % at 0712 hrs and 66 to 89 % at 1412 hrs. The winds run mainly from the ENE & ESE with a speed of 1.5 to 8.7 km/hour. Light rain may likely occur in the coming five days. Maximum & minimum temperature may range from 29.0 to 32.0 deg C & 24.0 to 25.0 deg C respectively. Wind will blow with a speed of 10.0-12.0 km/hr from the ESE direction. As per ERF products rainfall, maximum and minimum temperature will be normal from 1 - 7 September in the state.

### General Advisory:

In view of the second wave of Corona (Covid-19), the farmers are advised to follow the guidelines given by the Government of India, personal hygiene, use of masks, washing hands at appropriate intervals during agricultural operations. NDVI & SPI map received from the India Meteorological Department dated indicated that the NDVI ranged between 0.30-0.45 for U S Nagar i.e. agriculture vigor is moderate in the district. SPI map indicates the mildly dry conditions in the district for the past 4 weeks. Farmers are advised to download "Meghdoot" and "Damini" mobile apps on their smartphones from Google Play Store for free to get timely weather forecasts and district-wise Agromet advisory bulletins and lightning information.

### SMS Advisory:

Light rain may likely occur in the coming five days.

**Crop Specific Advisory:**

<b>Crop(Varieties)</b>	<b>Crop Specific Advisory</b>
RICE	In the early stage of paddy field, on the occurrence of stem borer, the field should be filled with 2 inches of water then spray, chlorantraniliprole 0.4G @10 kg/ha of or Fipronil 0.3G @ 25 kg/ha within 50 days of transplanting
MAIZE	If more than 5% of maize plants are affected by borer, then spray Coragen @ 150 ml per hectare. Spraying of chemicals should be done in clear weather.

**Horticulture Specific Advisory:**

<b>Horticulture(Varieties)</b>	<b>Horticulture Specific Advisory</b>
CAULIFLOWER	Browning or brown rot of cauliflower is due to boron deficiency; it appears on water-soaked areas and later changes into rusty brown. Spray 1kg of Borax in 500 liters of water 30 days after planting. Use the chemicals in clear weather conditions.
LEMON	To prevent the citrus canker disease in lemon, first of all, the diseased twigs should be cut and sprayed with blue copper 3 g / liter medicine after giving proper shape to the plants. Keep in mind that the diseased twigs should be burnt far away from the plants.
CAPSICUM	Sowing of mid-duration varieties of capsicum and tomato should be completed.

**Live Stock Specific Advisory:**

<b>Live Stock(Varieties)</b>	<b>Live Stock Specific Advisory</b>
BUFFALO	Vaccinate the animals against FMD, if not already done so. Animals afflicted by FMD should be kept in a separate enclosure so that they do not infect the healthy ones. If FMD is prevalent in the area, do not let your animals come in contact with the infected ones.

**Fisheries Specific Advisory:**

<b>Fisheries(Varieties)</b>	<b>Fisheries Specific Advisory</b>
COMMON FISH	In this season, due to rain and cloud, there is a lack of oxygen and excess ammonia in the water, due to which the fish can die, so to avoid this, replace the old water with new water containing oxygen. Also, use toximar at the rate of 10 kg per acre in the pond.

**Poultry Specific Advisory:**

<b>Poultry(Varieties)</b>	<b>Poultry Specific Advisory</b>
CHICKEN	Protect the poultry farm from moisture, provide proper lighting and clean the dust and dirt of utensils.